

Research Citations from Get Moving

JAMA. 2005;293:2479-2486

Physical Activity and Survival After Breast Cancer Diagnosis

Michelle D. Holmes, MD, DrPH et al.

<http://jama.ama-assn.org/cgi/content/abstract/293/20/2479>

Cancer 2003;97:1746-57

The Health, Eating, Activity, and Lifestyle (HEAL) Study

Physical Activity Levels before and after a Diagnosis
of Breast Carcinoma

Irwin. et al.

<http://appliedresearch.cancer.gov/surveys/heal/>

PAL Trial-Lymphedema

Contemp Clin Trials. 2009 May;30(3):233-45. Epub 2009 Jan 8.

Physical Activity and Lymphedema (the PAL trial): assessing the safety of progressive strength training in breast cancer survivors.

KH,

<http://www.ncbi.nlm.nih.gov/pubmed/19171204>

BMJ. 2009 Oct 13;339:b3410. doi: 10.1136/bmj.b3410.

of a multimodal high intensity exercise intervention in cancer patients undergoing chemotherapy: randomised controlled trial.

Adamsen L et al.

http://www.bmj.com/cgi/content/abstract/339/oct13_1/b3410

Feasibility of a 6-Month Diet, Exercise, and Behavior Modification Intervention for Post-Menopausal Breast Cancer Survivors

http://www.abstracts2view.com/sabcs09/view.php?nu=SABCS09L_856